

Total number of printed pages-32.

3 (Sem-5/CBCS) MAT HE 1/HE 2/HE 3

2022

MATHEMATICS

(Honours Elective)

Answer the Questions from any one Option.

OPTION-A


Paper : MAT-HE-5016

(Number Theory)

DSE (H)-1

Full Marks : 80

Time : Three hours



The figures in the margin indicate full marks for the questions.


PART-A

1. Choose the correct option in each of the following questions : **(any ten)** $1 \times 10 = 10$
- (i) Number of integers which are less than and co-prime to 108 is
- (a) 18
- (b) 17

Contd.

- (c) 15
(d) 36
- (ii) The number of positive divisors of a perfect square number is
- (a) odd
(b) even
(c) prime
(d) Can't say
- (iii) If $100! \equiv x \pmod{101}$, then x is
- (a) 99
(b) 100
(c) 101
(d) None of the above
- (iv) The solution of pair of linear congruences $2x \equiv 1 \pmod{5}$ and $x \equiv 4 \pmod{3}$ is
- (a) $x \equiv 13 \pmod{5}$
(b) $x \equiv 28 \pmod{5}$
(c) $x \equiv 13 \pmod{15}$
(d) $x \equiv 13 \pmod{3}$

- (v) If $a = qb$ for some integer q and $a, b \neq 0$, then
- (a) b divides a
(b) a divides b
(c) $a = b$
(d) None of the above
- (vi) If a and b are any two integers, then there exists some integers x and y such that
- (a) $\gcd(a, b) = ax + by$
(b) $\gcd(a, b) = ax - by$
(c) $\gcd(a, b) = ax^n + by^m$
(d) $\gcd(a, b) = (ax + by)^n$
- (vii) The linear diophantine equation $ax + by = c$ with $d = \gcd(a, b)$ has a solution in integers if and only if
- (a) $d \mid c$
(b) $c \mid d$
(c) $d \mid (ax + by)$
(d) Both (a) and (c)



(viii) If a positive integer n divides the difference of two integers a and b , then

- (a) $a \equiv b \pmod{n}$
- (b) $a = b \pmod{n}$
- (c) $a \equiv n \pmod{b}$
- (d) None of the above

(ix) The set of integers such that every integer is congruent modulo m to exactly one integer of the set is called _____ modulo m . *(Fill in the blank)*

- (a) Reduced residue system
- (b) Complete residue system
- (c) Elementary residue system
- (d) None of the above

(x) Which of the following statement is false?

- (a) There is no pattern in prime numbers
- (b) No formulae for finding prime numbers
- (c) Both (a) and (b)
- (d) None of the above

(xi) The reduced residue system is _____ of complete residue system.


- (a) compliment
- (b) subset
- (c) not a subset
- (d) Both (a) and (c)

(xii) The unit place digit of 137^{93} is

- (a) 7
- (b) 9
- (c) 3
- (d) 1

(xiii) Euler phi-function of a prime number p is

- (a) p
- (b) $p-1$
- (c) $p/2 - 1$
- (d) None of the above



(xiv) Which theorem states that "if p is prime, then $(p-1)! \equiv -1 \pmod{p}$ " ?

- (a) Dirichlet's theorem
- (b) Wilson's theorem
- (c) Euler's theorem
- (d) Fermat's little theorem

(xv) Let p be an odd prime. Then $x^2 \equiv -1 \pmod{p}$ has a solution if p is of the form

- (a) $4k+1$
- (b) $4k$
- (c) $4k+3$
- (d) None of the above

(xvi) Let m be a positive integer. Two integers a and b are congruent modulo m if and only if

- (a) $m \mid (a-b)$
- (b) $m \mid (a+b)$
- (c) $m \mid (ab)$
- (d) Both (b) and (c)

(xvii) If $ac \equiv bc \pmod{m}$ and $d = \gcd(m, c)$

(a) $a \equiv b \pmod{\frac{m}{d}}$

(b) $a \equiv c \pmod{\frac{m}{d}}$

(c) $a \equiv m \pmod{b}$

(d) $a \equiv m \pmod{\frac{b}{a}}$

(xviii) If a is a whole number and p is a prime number, then according to Fermat's theorem

(a) $a^p - a$ is divisible by p

(b) $a^p - 1$ is divisible by p

(c) $a^{p-1} - 1$ is divisible by p

(d) $a^{p-1} - a$ is divisible by p

2. Answer **any five** questions : $2 \times 5 = 10$

(a) Find last two digits of 3^{100} in its decimal expansion.

- (b) If p and q are positive integers such that $\gcd(p, q) = 1$, then show that $\gcd(a + b, a - b) = 1$ or 2 .
- (c) Find the solution of the following linear Diophantine equation $8x - 10y = 42$.
- (d) If p and q are any two real numbers, then prove that $[p] + [q] \leq [p + q]$ (where $[x]$ denotes the greatest integer less or equal to x).
- (e) If m and n are integers such that $(m, n) = 1$, then $\phi(mn) = \phi(m)\phi(n)$.
- (f) Find (7056) .
- (g) If $a \equiv b \pmod{n}$ and $m \mid n$, then show that $a \equiv b \pmod{m}$.
- (h) List all primitive roots modulo 7 .
- (i) If $n = p_1^{k_1} p_2^{k_2} \dots p_m^{k_m}$ is the prime factorization of $n > 1$, then prove that $\tau(n) = (k_1 + 1)(k_2 + 1) \dots (k_m + 1)$.
- (j) Evaluate the exponent of 7 in $1000!$

3. Answer **any four** questions : $5 \times 4 = 20$

- (a) If p is a prime, then prove that $\phi(p!) = (p-1)\phi((p-1)!)$
- (b) Show that, the set of integers $\{1, 5, 7, 11\}$ is a reduced residue system (RRS) modulo 12 .
- (c) Solve the following simultaneous congruence :

$$x \equiv 2 \pmod{3}$$

$$x \equiv 2 \pmod{2}$$

$$x \equiv 3 \pmod{5}$$
- (d) For $n = p^k$, p is a prime, prove that $n = \sum_{d \mid n} \phi(d)$, where $\sum_{d \mid n}$ denotes the sum over all positive divisors of n .
- (e) If p_n is the n^{th} prime, then show that $\frac{1}{p_1} + \frac{1}{p_2} + \dots + \frac{1}{p_n}$ is not an integer.
- (f) Let n be any integer > 2 . Then $\phi(n)$ is even.
- (g) Show that if $a_1, a_2, \dots, a_{\phi(m)}$ is a RRS modulo m , where m is a positive integer with $m \neq 2$, then $a_1 + a_2 + \dots + a_{\phi(m)} \equiv 0 \pmod{m}$.



- (h) Show that $10! + 1$ is divisible by 11.

PART-B

Answer **any four** of the following questions :

$$10 \times 4 = 40$$

4. (a) If $a, b \neq 0$ and c be any three integers and $d = \gcd(a, b)$. Then show that $ax + by = c$ has a solution iff $d \mid c$.

Furthermore, show that if x_0 and y_0 is a particular solution of $ax + by = c$, then any other solution of the equation is $x' = x_0 - \frac{b}{d}t$ and $y' = y_0 + \frac{a}{d}t$, t is an integer.

7

- (b) Find the general solution of $10x - 8y = 42$; $x, y \in \mathbb{Z}$

3

5. (a) Show that an odd prime p can be represented as sum of two squares iff $p \equiv 1 \pmod{4}$.

7

- (b) Find all positive solutions of $x^2 + y^2 = z^2$, where $0 < z < 30$.

3

6. (a) Prove that the quadratic congruence $x^2 + 1 \equiv 0 \pmod{p}$, where p is an odd prime, has a solution iff $p \equiv 1 \pmod{4}$.

5

- (b) If p is prime and a is an integer not divisible by p , prove that $a^{p-1} \equiv 1 \pmod{p}$.

5

7. State and prove Chinese remainder theorem. Also find all integers that leave a remainder of 4 when divided by 11 and leaves a remainder of 3 when divided by 17.

8. (a) For each positive integer $n \geq 1$, show

$$\text{that } \sum_{d|n} \mu(d) = \begin{cases} 1, & \text{if } n = 1 \\ 0, & \text{if } n > 1 \end{cases}$$


5

- (b) If k denotes the number of distinct prime factors of positive integer n . Prove that $\sum_{d|n} |\mu(d)| = 2^k$

5

9. (a) If p is a prime, prove that $\phi(p^k) = p^k - p^{k-1}$, for any positive integer k . For $n > 2$, show that $\phi(n)$ is an even integer.

3+2=5



- (b) State Mobius inversion formula. If the integer $n > 1$ has the prime factorization. If $n = p_1^{k_1} p_2^{k_2} \dots p_s^{k_s}$, then prove that

$$\sum_{d|n} \mu(d) \sigma(d) = (-1)^s p_1 p_2 \dots p_s. \quad 5$$

10. If x be any real number. Then show that
 $1+3+3+3=10$

(a) $[x] \leq x < [x] + 1$

(b) $[x+m] = [x] + m$, m is any integer

(c) $[x] + [-x] = \begin{cases} 0, & \text{if } x \text{ is an integer} \\ -1, & \text{otherwise} \end{cases}$

(d) $\left[\frac{[x]}{m}\right] = \left[\frac{x}{m}\right]$, if m is a positive integer

11. (a) If $a_1, a_2, \dots, a_m$ is a complete residue system modulo m , and if k is a positive integer with $(k, m) = 1$ then $ka_1 + b, ka_2 + b, \dots, ka_m + b$, is a complete residue system modulo m for any integer b .

- (b) Examine whether the following set forms a complete residue system or a reduced residue system :

$$\{-3, 14, 3, 12, 37, 56, -1\} \pmod{7} \quad 5$$

12. (a) If $n \geq 1$ is an integer then show that

$$\prod_{d|n} d = n^{\frac{\tau(n)}{2}} \quad 3$$


- (b) If f and g are two arithmetic functions, then show that the following conditions are equivalent : 7

(i) $f(n) = \sum_{d|n} g(d)$

(ii) $g(n) = \sum_{d|n} \mu(d) f\left(\frac{n}{d}\right) = \sum_{d|n} \mu\left(\frac{n}{d}\right) f(d)$

13. (a) If n is a positive integer with $n \geq 2$, such that $(n-1)! + 1 \equiv 0 \pmod{n}$, then show that n is prime. 5

- (b) Show that if p is an odd prime, then $2(p-3)! \equiv -1 \pmod{p}$. 5



OPTION-B

Paper : MAT-HE-5026

(Mechanics)

1. Answer the following questions : **(any ten)**
 $1 \times 10 = 10$

- (i) If a system of coplanar forces is in equilibrium, then what is the algebraic sum of the moment of the forces about any point in the plane ?
- (ii) What is the resultant of the like parallel forces $P_1, P_2, P_3, \dots$ acting on a body ?
- (iii) If a particle moves under the action of a conservative system of forces, then what is the sum of its KE and PE ?
- (iv) Define limiting equilibrium.
- (v) Define the centre of gravity of a body.
- (vi) Under what conditions the effect of a couple is not altered if it is transformed to a parallel plane ?
- (vii) Write down the radial and cross-radial components of velocities of a particle moving on a plane curve at any point (r, θ) on it.

- (viii) What is the resultant of a couple and a force in the same plane ?
- (ix) What is dynamical friction ?
- (x) What do you mean by terminal velocity ?
- (xi) Define coefficient of friction.
- (xii) What is the position of the point of action of the resultant of two equal like parallel forces acting on a rigid body ?
- (xiii) What is the whole effect of a couple acting on a body ?
- (xiv) Define simple harmonic motion.
- (xv) What is the centre of gravity of a triangular lamina ?
- (xvi) Define limiting friction.
- (xvii) State the principle of conservation of energy.
- (xviii) A particle moves on a straight line towards a fixed point O with an acceleration proportional to its distance from O . If x is the distance of the particle at time t from O , then write down its equation of motion.

2. Answer **any five** questions of the following :
2×5=10

- (a) Write the laws of static friction.
- (b) A particle moves in a circle of radius r with a speed v . Prove that its angular velocity is $\frac{v}{r}$.
- (c) What are the general conditions of equilibrium of any system of coplanar forces ?
- (d) The law of motion in a straight line is $s = \frac{1}{2}vt$. Prove that the acceleration is constant.
- (e) Find the greatest and least resultant of two forces acting at a point whose magnitudes are P and Q respectively.
- (f) Find the centre of gravity of an arc of a plane curve $y = f(x)$.
- (g) State Hooke's law.
- (h) Show that impulse of a force is equal to the momentum generated by the force in the given time.

- (i) Write the expression for the component of velocity and acceleration along radial and cross radial direction in a motion of a particle in a plane curve.
- (j) The speed v of a particle moving along x -axis is given by the relation $v^2 = n^2(8bx - x^2 - 12b^2)$. Prove that the motion is Simple Harmonic.

3. Answer **any four** questions of the following :
5×4=20

- (a) The greatest and least resultants that two forces acting at a point can have magnitude P and Q respectively. Show that when they act at an angle α their resultant is $\sqrt{P^2 \cos^2 \frac{\alpha}{2} + Q^2 \sin^2 \frac{\alpha}{2}}$.
- (b) I is the in centre of the triangle ABC . If three forces $\vec{P}, \vec{Q}, \vec{R}$ acting at I along $\vec{IA}, \vec{IB}, \vec{IC}$ are in equilibrium, prove that

$$\frac{P}{\sqrt{a(b+c-a)}} = \frac{Q}{\sqrt{b(c+a-b)}} = \frac{R}{\sqrt{c(a+b-c)}}$$



- (c) Show that the resultant of three equal like parallel forces acting at the three vertices of a triangle passes through the centroid of the triangle.
- (d) Prove that any system of coplanar forces acting on a rigid body can ultimately be reduced to a single force acting at any arbitrarily chosen point in the plane, together with a couple.
- (e) Show that the sum of the Kinetic energy and Potential energy is constant throughout the motion when a particle of mass m falls from rest at a height h above ground.
- (f) A point moves along a circle with constant speed. Find its angular velocity and acceleration about any point of the circle.
- (g) Show that the work done against tension in stretching a light elastic string is equal to the product of its extension and the mean of the initial and final tension.
- (h) A particle starts with velocity u and moves under retardation μ times of the distance. Show that the distance it

travels before it comes to rest is $\frac{u}{\sqrt{\mu}}$.

4. Answer **any four** questions of the following :

$$10 \times 4 = 40$$

- (a) Forces P , Q and R act along the sides BC , CA and AB of a triangle ABC and forces P' , Q' and R' act along OA , OB and OC , where O is the centre of the circumscribed circle, prove that

$$(i) \quad P \cos A + Q \cos B + R \cos C = 0$$

$$(ii) \quad \frac{PP'}{a} + \frac{QQ'}{b} + \frac{RR'}{c} = 0$$

- (b) State and prove Lami's theorem. Forces P , Q and R acting along OA , OB and OC , where O is the circumcentre of triangle ABC , are in equilibrium. Show that

$$\frac{P}{a^2(b^2 + c^2 - a^2)} = \frac{Q}{b^2(c^2 + a^2 - b^2)} = \frac{R}{c^2(a^2 + b^2 - c^2)}$$

- (c) (i) Find the centre of gravity of a uniform arc of the circle $x^2 + y^2 = a^2$ in the positive quadrant.



- (ii) Find the centre of gravity of the arc of the asteroid $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$ lying in the first quadrant.
- (d) A particle moves in a straight line under an attraction towards a fixed point on the line varying inversely as the square of the distance from the fixed point. Investigate the motion.
- (e) A particle moves in a straight line OA starting from the rest at A and moving with an acceleration which is directed towards O and varies as the distance from O . Discuss the motion of the particle. Hence define Simple Harmonic Motion and time period of the motion.
- (f) Find the component of acceleration of a point moving in a plane curve along the initial line and the radius vector. Also find the component of acceleration perpendicular to initial line and perpendicular to radius vector.
- (g) A particle is falling under gravity in a medium whose resistance varies as the velocity. Find the distance and velocity at any time t . Also find the terminal velocity of the particle.

- (h) The velocity component of a particle along and perpendicular to the radius vector from λr and $\mu\theta$. Find the path and show that radial and transverse component of acceleration are

$$\lambda^2 r - \frac{\mu^2 \theta^2}{r} \text{ and } \mu\theta \left(\lambda + \frac{\mu}{r} \right).$$

- (i) Find the component of acceleration of a point moving in a plane curve along the initial line and the radius vector. Also find the component of acceleration perpendicular to initial line and perpendicular to radius vector.
- (j) A particle moves in a straight line under an attraction towards a fixed point on the line varying inversely as the square of the distance from the fixed point. Investigate the motion.



OPTION-C

Paper : MAT-HE-5036

(Probability and Statistics)

1. Answer **any ten** questions from the following :

$$1 \times 10 = 10$$

- (a) If A and B are mutually exclusive what will be the modified statement of

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

- (b) Define probability density function for a continuous random variable.

- (c) A random variable X can take all non-negative integral values, and the probability that X takes the value r is

$$P(X=r) = A\alpha^r \quad (0 < \alpha < 1). \text{ Find } P(X=0).$$

- (d) If X and Y are two random variables and $\text{var}(X-Y) \neq \text{var}(X) - \text{var}(Y)$ then what is the relation between X and Y .

- (e) Test the velocity of the following probability distribution :

x	-1	0	1
$P(x)$	0.4	0.4	0.3

- (f) Define Negative Binomial distribution for a random variable X with parameter r .

- (g) What are the relations between mean, median and mode of a normal distribution ?

- (h) Write the equation of the line of regression of x on y .

- (i) What is the variance of the mean of a random sample ?

- (j) Define moment generating function of a random variable X about origin.

- (k) What are the limits for correlation coefficients ?

- (l) For a Bernoulli random variable X with $P(X=0) = 1-P$ and $P(X=1) = P$ write $E(X)$ and $V(X)$ in terms of P .

- (m) If X is a random variable with mean μ and variance σ^2 , then for any positive number k , find Chebychev's inequality.

- (n) A continuous random variable X follow the probability law $f(x) = Ax^2$, $0 \leq x \leq 1$. Determine A .



(o) If X and Y are two random variables then find $cov(x, y)$.

(p) If a is constant then find $E(a)$ and $var(a)$.

(q) If X and Y are two independent. Poisson variates, then XY is a _____ variate.
(Fill in the blank)

(r) If a non-negative real valued function f is the probability density function of some continuous random variable, then

what is the value of $\int_{-\alpha}^{\alpha} f(x) dx$?

2. Answer the following questions : **(any five)**
 $2 \times 5 = 10$

(a) If A and B are independent events, then show that A and B are also independent.

(b) If X have the p.m.f

$$f(x) = \frac{x}{10}, x = 1, 2, 3, 4$$

then find $E(X^2)$

(c) With usual notation for a binomial variate X , given that

$$9P(x=4) = P(x=2) \text{ when } n=6.$$

Find the value of p and q .

(d) If X is a continuous random variable whose probability density function is given by

$$f(x) = \begin{cases} \frac{3}{8}(4x - 2x^2) & 0 < x < 2 \\ 0, & \text{otherwise} \end{cases}$$

then find $P\{X > 1\}$.

(e) Show that for a normal standard variate z , $E(z) = 0$ and $V(z) = 1$.

(f) The number of items produced in a factory during a week is a random variable with mean 50. If the variance of a week's production is 25, then what is the probability that this week's production will be between 40 and 60.

(g) Define probability mass function and probability density function for a random variable X .



- (h) If X is a random Poisson variate with parameter λ , then show that

$$P(X \geq n) - P(X \geq n+1) = \frac{e^{-\lambda} \lambda^n}{n}$$

- (i) If $M_X(t)$ is a moment generating function of a random variable X with parameter t then show that

$$M_{cX}(t) = M_X(ct), c \text{ is a constant.}$$

- (j) If X and Y are independent random variables with characteristic functions $\phi_X(w)$ and $\phi_Y(w)$ respectively then show that

$$\phi_{X+Y}(w) = \phi_X(w)\phi_Y(w)$$

3. Answer **any four** questions from the following : 5×4=20

- (a) If X is a discrete random variable having probability mass function 2+2+1=5

mass point	0	1	2	3	4	5	6	7
$p(X=x)$	0	k	$2k$	$3k$	$4k$	k^2	$2k^2$	$7k^2 + k$

Determine :

- (i) k
(ii) $p(X < 6)$ and
(iii) $p(X \geq 6)$



- (b) If X and Y are two independent random variables then show that

$$\text{var}(X+Y) = \text{var}(X) + \text{var}(Y)$$

- (c) If the probability that an individual will suffer a bad reaction from injective of a given serum is 0.001 determine the probability that out of 2000 individuals

- (i) exactly 3,
(ii) more than 2 individual

will suffer a bad reaction. 2+3=5

- (d) Two random variables X and Y are jointly distributed as follows :

$$f(x, y) = \frac{2}{\pi} (1 - x^2 - y^2), 0 < x^2 + y^2 < 1$$

Find the marginal distribution of X .


- (e) State and prove weak law of large numbers.

- (f) If X and Y are independent random variables having common density function

$$f(x) = e^{-x}, x > 0$$

0, otherwise

Find the density function of the random variable X/Y .



- (g) If X and Y are independent Poisson variates such that

$$P(x=1) = P(x=2) \text{ and}$$

$$P(y=2) = P(y=3)$$

Find the variance of $x - 2y$.

- (h) Prove that regression coefficients are independent of the change of origin but not of scale.

4. Answer **any four** from the following questions : 10×4=40

- (a) (i) What is meant by partition of a sample space S ? If $H_i (i=1, 2, \dots, n)$ is a partition of the sample space S , then for any event A , prove that

$$P(H_i/A) = \frac{P(H_i)P(A/H_i)}{\sum_{i=1}^n P(H_i)P(A/H_i)} \quad 5$$

- (ii) If X is a random variable with the following probability distribution :

$$x: -3 \quad 6 \quad 9$$

$$P(X=x): 1/6 \quad 1/2 \quad 1/3$$

Find $E(X)$, $E(X^2)$ and $var(X)$ 5

- (b) Two random variables X and Y have the following joint probability density function : 2+2+3+3=10



$$f(x, y) = \begin{cases} 2 - x - y, & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Find (i) marginal probability density function of X and Y (ii) conditional density function (iii) $var(X)$ and $var(Y)$ (iv) co-variance between X and Y .

- (c) (i) Let X be a random variable with mean μ and variance r^2 . Show that $E(x-b)^2$ as a function of b is minimum when $b = \mu$. 5

- (ii) A bag contains 5 balls and it is known how many of these are white. Two balls are drawn and are found to be white. What is the probability that all are white? 5

- (d) (i) If X is a random variable then prove that 5
- $$var(X) = var[E(X/Y)] + E[var(X/Y)]$$



- (ii) Find the probability that in a family of 4 children there will be
(a) at least one boy (b) at least one boy and at least one girl. 5

- (e) What are the chief characteristics of the normal distribution and normal curve? 5

- (f) (i) Show that mean and variance of a Poisson distribution are equal. 5

- (ii) Determine the binomial distribution for which the mean is 4 and variance is 3 and find its mode. 5

- (g) (i) Prove that independent variables are uncorrelated. With the help of an example show that the converse is not true. 5

- (ii) Find the angle between the two lines of regression 5

$$y - \bar{y} = \frac{r\sigma_y}{\sigma_x} (x - \bar{x})$$

$$\text{and } x - \bar{x} = r \frac{\sigma_x}{\sigma_y} (y - \bar{y})$$

- (h) (i) A function $f(x)$ of x is defined as follows :

$$f(x) = 0 \text{ for } x < 2$$

$$= \frac{1}{18} (3 + 2x) \text{ for } 2 \leq x \leq 4$$

$$= 0, \text{ for } x > 4$$

Show that it is a density function. Also find the probability that a variate with this density will lie in the interval $2 \leq x \leq 3$. 5

- (ii) A random variable X can assume values 1 and -1 with probability $\frac{1}{2}$ each. Find



- (i) moment generating function,
(ii) characteristics function. 5

- (i) (i) Find the median of a normal distribution. 5

- (ii) A random variable X has density function given by 5

$$f(x) = \begin{cases} 2e^{-2x}, & x \geq 0 \\ 0, & x < 0 \end{cases}$$

Find (i) mean with the help of m.g.f. (ii) $P[|x - \mu| > 1]$.





- (j) (i) The diameter say x , of an electric cable is assumed to be continuous random variable with p.d.f

$$f(x) = 6x(1-x), 0 \leq x \leq 1$$

- (a) Check that the above is a p.d.f,

- (b) Determine the value of k such that $P(X < K) = P(X > K)$ 5

- (ii) If 3% of electric bulbs manufactured by a company are defective, using Poisson's distribution find the probability that in a sample of 100 bulbs exactly 5 bulbs are defective 5

$$[\text{Given } e^{-3} = 0.04979]$$

